

Please click price below and you will be redirected to our pharmacy.

## Isordil



**Generic Isordil** is used for preventing symptoms of angina (chest pain) caused by heart disease. This medicine is used alone or with other medicines. This medicine is NOT intended for the immediate relief of acute attacks of angina.

Active ingredient: [Isosorbide Dinitrate](#)

**Brand(s):**Isordil-10  
**Manufacturer:** Ipca Laboratories  
**Disease(s):**Angina

### Isordil 10mg

	Per pill	Per Pack	Order
30 tablet	\$ 1.25	<b>\$ 37.50</b>	<a href="#">Add to cart</a>
60 tablet	\$ 1.20	<b>\$ 72.00</b> <del>\$75.00</del>	<a href="#">Add to cart</a>
90 tablet	\$ 1.10	<b>\$ 99.00</b> <del>\$112.50</del>	<a href="#">Add to cart</a>
120 tablet	\$ 1.00	<b>\$ 120.00</b> <del>\$150.00</del>	<a href="#">Add to cart</a>
180 tablet	\$ 0.90	<b>\$ 162.00</b> <del>\$225.00</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
360 tablet	\$ 0.80	<b>\$ 288.00</b> <del>\$450.00</del>	<a href="#">Add to cart</a>
Free AirMail shipping			

## Isosorbide Dinitrate tablet

### What is this medicine?

ISOSORBIDE DINITRATE is a type of vasodilator. It relaxes blood vessels, increasing the blood and oxygen supply to your heart. This medicine is used to prevent chest pain caused by angina. It will not help to stop an episode of chest pain.

### What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions:

- previous heart attack or heart failure
- an unusual or allergic reaction to isosorbide dinitrate, nitrates, other medicines, foods, dyes, or preservatives
- pregnant or trying to get pregnant
- breast-feeding

### How should I use this medicine?

Take this medicine by mouth with a glass of water. Take this medicine on an empty stomach, at least 30 minutes before or 2 hours after food. Do not take with food. Take your medicine at regular intervals. Do not take your medicine more often than directed. Do not stop taking this medicine suddenly or your symptoms may get worse. Ask your doctor or health care professional how to gradually reduce the dose.

Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.

**Overdosage:** If you think you have taken too much of this medicine

