

Please click price below and you will be redirected to our pharmacy.

## Samsca



Tolvaptan reduces the level of a hormone that regulates the balance of water and salt (sodium) in the body. High levels of this hormone can cause an imbalance that results in low sodium levels and fluid retention. Tolvaptan is used to treat hyponatremia (low levels of sodium in your blood) in people with heart failure, and certain hormonal imbalances. Tolvaptan improves urine flow without causing the body to lose too much sodium as you urinate.

Active ingredient: [Tolvaptan](#)

**Brand(s):**Resodim  
**Manufacturer:** Lupin  
**Disease(s):**Hyponatremia

### Samsca 15mg

	Per pill	Per Pack	Order
20 tablet	\$ 15.00	<b>\$ 299.95</b>	<a href="#">Add to cart</a>
Free AirMail shipping			
40 tablet	\$ 13.00	<b>\$ 519.95</b> <del>\$599.90</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
60 tablet	\$ 12.00	<b>\$ 719.95</b> <del>\$899.85</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
120 tablet	\$ 10.00	<b>\$ 1199.95</b> <del>\$1799.70</del>	<a href="#">Add to cart</a>
Free AirMail shipping			

## Tolvaptan tablets

### What is tolvaptan?

Tolvaptan reduces the level of a hormone that regulates the balance of water and salt (sodium) in the body. High levels of this hormone can cause an imbalance that results in low sodium levels and fluid retention. Tolvaptan is used to treat hyponatremia (low levels of sodium in your blood) in people with heart failure, and certain hormonal imbalances. Tolvaptan improves urine flow without causing the body to lose too much sodium as you urinate.

Tolvaptan may also be used for purposes not listed in this medication guide.

### Important Information

You will need to be in a hospital under observation any time you start or restart treatment with tolvaptan.

Tell your doctor about all your current medicines and any you start or stop using. Many drugs can interact, and some drugs should not be used together.

Tell your caregivers at once if you have signs that your hyponatremia is being reversed too quickly: problems with speech or muscle control, trouble swallowing, trouble moving your arms and legs, confusion, or seizure.

While taking tolvaptan at home, call your doctor if you have signs of liver problems, such as: loss of appetite, upper stomach pain, dark urine, or jaundice (yellowing of the skin or eyes).

### Before taking this medicine

