

Please click price below and you will be redirected to our pharmacy.

## Ceclor CD



**Generic Ceclor** Modified-Release is used in the treatment of ear, nose, throat, respiratory tract, urinary tract, and skin infections caused by specific bacteria, including staph, strep, and E. coli. Uses include treatment of sore or strep throat, pneumonia, and tonsillitis. It is also used for flare-ups of chronic bronchitis.

Active ingredient: [Cefaclor](#)

**Brand(s):** Distaclor CD

**Manufacturer:** Baroque Pharmaceuticals Pvt. Ltd.

**Disease(s):** Infections / Pneumonia / Tonsillitis

### Ceclor CD 375mg

	Per pill	Per Pack	Order
10 tablet	\$ 7.46	<b>\$ 74.60</b>	<a href="#">Add to cart</a>
20 tablet	\$ 6.86	<b>\$ 137.20</b> <del>\$149.20</del>	<a href="#">Add to cart</a>
30 tablet	\$ 6.33	<b>\$ 189.80</b> <del>\$223.80</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
60 tablet	\$ 5.79	<b>\$ 347.60</b> <del>\$447.60</del>	<a href="#">Add to cart</a>

Free AirMail shipping

## Cefaclor Modified Release tablet

### What is this medicine?

CEFACLOR is an antibiotic. It is used to treat infections of the skin, respiratory tract, sinuses, ear, and urinary tract.

### What should my health care professional know before I receive this medicine?

They need to know if you have any of these conditions:

- bleeding problems
- diarrhea
- kidney disease
- stomach or intestine problems (especially colitis)
- an unusual or allergic reaction to cefaclor, other cephalosporin antibiotics, penicillin, penicillamine, other foods, dyes or preservatives
- pregnant or trying to get pregnant
- breast-feeding

### How should I use this medicine?

Take this medicine by mouth. Take it with food or within 1 hour after eating. Swallow the tablets whole. Do not chew, crush, or cut the tablets. Take your medicine at regular intervals. Do not take it more often than directed. Take all of your medicine as directed even if you think you are better. Do not skip doses or stop your medicine early.

Contact your pediatrician regarding the use of this medicine in children. Special care may be needed.

**Overdosage:** If you think you have taken too much of this medicine contact a poison control center or emergency room at once.

