

Ovol



Ovol allows gas bubbles in the stomach and intestines to come together more easily, which allows for easier passage of gas. Ovol is used to relieve painful pressure caused by excess gas in the stomach and intestines. Ovol is for use in babies, children, and adults.

Active ingredient: [Simethicone](#)

Brand(s):Gastica

Manufacturer: Mankind Pharma Limited

Disease(s):Gas / Spasms / Gripping / Tympanites / Flatulence / Bellyache / Stomach-ache / Tummy-ache

Ovol 15ml

| | Per bottle | Per Pack | Order |
|-----------|------------|---------------------------------------|-----------------------------|
| 1 bottles | \$ 18.95 | \$ 18.95 | Add to cart |
| 2 bottles | \$ 12.47 | \$ 24.95 \$37.90 | Add to cart |
| 3 bottles | \$ 10.98 | \$ 32.95 \$56.85 | Add to cart |
| 4 bottles | \$ 9.99 | \$ 39.95 \$75.80 | Add to cart |
| 5 bottles | \$ 8.99 | \$ 44.95 \$94.75 | Add to cart |

Simethicone drops

What is this medicine?

Simethicone allows gas bubbles in the stomach and intestines to come together more easily, which allows for easier passage of gas.

Simethicone is used to relieve painful pressure caused by excess gas in the stomach and intestines. Simethicone is for use in babies, children, and adults.

Simethicone may also be used for purposes not listed in this medication guide.

What should I tell my health care providers before I take this medicine?

You should not use this medication if you are allergic to simethicone.

Ask a doctor or pharmacist if it is safe for you to take this medicine if you are allergic to any drugs, or if you have any type of serious illness (especially one that affects your stomach or intestines).

Simethicone is not expected to harm an unborn baby.

It is not known whether simethicone passes into breast milk or if it could harm a nursing baby. Do not use this medication without medical advice if you are breast-feeding a baby.

The liquid form may contain phenylalanine. Talk to your doctor before using this form of simethicone if you have phenylketonuria (PKU).

How should I use this medicine?

Use exactly as directed on the label, or as prescribed by your doctor. Do not use in larger or smaller amounts or for longer than recommended. Do not take more of this medication than is directed. Simethicone works best if you take it after meals and at bedtime. The simethicone chewable tablet must be chewed before swallowing. Measure liquid medicine with a

