

## Zithromax Dispersible



**Generic** Zithromax(Azithromycin) dispersible is used to treat many different types of infections caused by bacteria, such as respiratory infections, skin infections, ear infections, and sexually transmitted diseases. In children, it is used to treat middle ear infection, pneumonia, tonsillitis, and strep throat.

Active ingredient: [Azithromycin](#)

**Brand(s):**AZEE DT

**Manufacturer:** Protec Pharma Ltd

**Disease(s):**Infections / Pneumonia / Tonsillitis

### Zithromax Dispersible 100mg

|           | Per pill | Per Pack                                | Order                       |
|-----------|----------|-----------------------------------------|-----------------------------|
| 12 tablet | \$ 4.91  | <b>\$ 58.95</b>                         | <a href="#">Add to cart</a> |
| 30 tablet | \$ 3.63  | <b>\$ 108.95</b><br><del>\$147.38</del> | <a href="#">Add to cart</a> |
| 60 tablet | \$ 3.27  | <b>\$ 195.95</b><br><del>\$294.75</del> | <a href="#">Add to cart</a> |

Free AirMail shipping

## Azithromycin tablet

### What is this medicine?

AZITHROMYCIN is a macrolide antibiotic that interferes with the growth of bacterial cells. It is used to treat bacterial infections in many different parts of the body. Azithromycin also treats sexually transmitted vaginal or urinary tract infections caused by chlamydia. It will not work for colds, flu, or other virus infections.

### What should my health care professional know before I take this medicine?

They need to know if you have any of these conditions;;

- kidney disease; liver disease
- pneumonia
- stomach problems (especially colitis)
- other chronic illness; an unusual or allergic reaction to azithromycin
- other macrolide antibiotics (such as erythromycin), foods, dyes, or preservatives
- pregnant or trying to get pregnant
- breast-feeding

### How should I take this medicine?

Azithromycin tablets can be taken with or without food. Take your doses at regular intervals. Do not take your medicine more often than directed. Finish the full course prescribed by your prescriber or health care professional even if you think your condition is better. Do not stop taking except on your prescriber's advice. Contact your pediatrician or health care professional regarding the use of this medicine in children. Special care may be needed. Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once. NOTE: This medicine is only for you. Do not share this medicine with others.

### What if I miss a dose?

