

## Procoralan



**Procoralan** works by affecting your heart's electrical activity in order to slow the heart rate. Ivabradine is used in certain people with chronic heart failure, to help lower the risk of needing to be hospitalized when symptoms get worse. Ivabradine is not for use if you already have heart failure symptoms that recently got worse before starting this medicine. Ivabradine may also be used for purposes not listed in this medication guide.

Active ingredient: [Ivabradine](#)

**Brand(s):**Ivabeat

**Manufacturer:** Cipla Limited

**Disease(s):**Chronic Heart Failure

### Procoralan 5mg

	Per pill	Per Pack	Order
10 tablet	\$ 2.50	<b>\$ 24.95</b>	<a href="#">Add to cart</a>
20 tablet	\$ 2.00	<b>\$ 39.95</b> <del>\$49.90</del>	<a href="#">Add to cart</a>
30 tablet	\$ 1.67	<b>\$ 49.95</b> <del>\$74.85</del>	<a href="#">Add to cart</a>
60 tablet	\$ 1.50	<b>\$ 89.95</b> <del>\$149.70</del>	<a href="#">Add to cart</a>
90 tablet	\$ 1.44	<b>\$ 129.95</b> <del>\$224.55</del>	<a href="#">Add to cart</a>
120 tablet	\$ 1.37	<b>\$ 164.95</b> <del>\$299.40</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
180 tablet	\$ 1.33	<b>\$ 239.95</b> <del>\$449.10</del>	<a href="#">Add to cart</a>
Free AirMail shipping			

## Ivabradine tablet

### What is this medicine?

IVABRDINE works by affecting your heart's electrical activity in order to slow the heart rate. Ivabradine is used in certain people with chronic heart failure, to help lower the risk of needing to be hospitalized when symptoms get worse. Ivabradine is not for use if you already have heart failure symptoms that recently got worse before starting this medicine. Ivabradine may also be used for purposes not listed in this medication guide.

### What should I tell my health care providers before I take this medicine?

You should not use ivabradine if you are allergic to it, or if you have:

- a serious heart condition such as "sick sinus syndrome" or "AV block" (unless you have a pacemaker);
- heart failure symptoms that recently got worse;
- severe liver disease;
- if your resting heart rate is less than 60 beats per minute;
- if your blood pressure is lower than 90 over 50; or
- if you depend on a pacemaker to regulate your heart rate.

Some medicines can cause unwanted or dangerous effects when used with ivabradine. Tell your doctor about all the medicines you are taking, including over-the-counter medicines, vitamins, and herbal products.

