

## Droxia



Generic **Droxia** is used for treating skin cancer, cancer of the ovary or chronic myelocytic leukemia that is recurrent, has spread or cannot be helped with surgery. It may also be used with radiation to control skin cancers of the head and neck.

Active ingredient: [Hydroxyurea](#)

**Brand(s):**Hydrea

**Manufacturer:** Baramhaj Chemicals

**Disease(s):**Leukemia / Skin Cancer / Ovarian Cancer / Sickle Cell Anemia

### Droxia 500mg

	Per pill	Per Pack	Order
30 tablet	\$ 4.04	<b>\$ 121.20</b>	<a href="#">Add to cart</a>
60 tablet	\$ 3.34	<b>\$ 200.40</b> <del>\$242.40</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
90 tablet	\$ 3.11	<b>\$ 279.60</b> <del>\$363.60</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
120 tablet	\$ 2.99	<b>\$ 358.80</b> <del>\$484.80</del>	<a href="#">Add to cart</a>
Free AirMail shipping			

## Hydroxyurea capsule

### What is this medicine?

HYDROXYUREA is a chemotherapy drug. It slows the growth of cancer cells. This medicine is used to treat certain leukemias, skin cancer, head and neck cancer, and advanced ovarian cancer. It is also used to control the painful crises of sickle cell anemia.

### What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions:

- immune system problems
- infection (especially a virus infection such as chickenpox, cold sores, or herpes)
- kidney disease
- low blood counts, like low white cell, platelet, or red cell counts
- previous or ongoing radiation therapy
- an unusual or allergic reaction to hydroxyurea, other chemotherapy, other medicines, foods, dyes, or preservatives
- pregnant or trying to get pregnant
- breast-feeding

### How should I use this medicine?

Take this medicine by mouth with a glass of water. Take your medicine at regular intervals. Do not take it more often than directed. Do not stop taking except on your doctor's advice.

People who are not taking this medicine should not be exposed to it. Wash your hands before and after handling your bottle or medicine. Caregivers should wear disposable gloves if they must touch the bottle or medicine. Clean up any medicine powder that spills with a damp

