

Please click price below and you will be redirected to our pharmacy.

## Vastarel



**Vastarel** is an antianginal drug. Normalizes the energy metabolism of cells undergoing hypoxia or ischemia. Supports energy metabolism of the heart and neurosensory organs. Used for the prevention of angina attacks in coronary heart disease, as well as in Meniere's disease, tinnitus, dizziness caused by a cerebrovascular accident.

Active ingredient: [Trimetazidine](#)

**Brand(s):** Flavedon

**Manufacturer:** Serdia

**Disease(s):** Angina Attacks / Coronary Heart Disease / Meniere's Disease / Tinnitus / Dizziness /

### Vastarel 20mg

	Per capsule	Per Pack	Order
30 capsules	\$ 1.50	<b>\$ 44.95</b>	<a href="#">Add to cart</a>
60 capsules	\$ 1.40	<b>\$ 83.95</b> <del>\$89.90</del>	<a href="#">Add to cart</a>
90 capsules	\$ 1.20	<b>\$ 107.95</b> <del>\$134.85</del>	<a href="#">Add to cart</a>
120 capsules	\$ 1.10	<b>\$ 131.95</b> <del>\$179.80</del>	<a href="#">Add to cart</a>
180 capsules	\$ 1.00	<b>\$ 179.95</b> <del>\$269.70</del>	<a href="#">Add to cart</a>

Free AirMail shipping

### Vastarel 60mg

	Per capsule	Per Pack	Order
30 capsules	\$ 3.50	<b>\$ 104.95</b>	<a href="#">Add to cart</a>
60 capsules	\$ 3.33	<b>\$ 199.95</b> <del>\$209.90</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
90 capsules	\$ 3.11	<b>\$ 279.95</b> <del>\$314.85</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
120 capsules	\$ 2.96	<b>\$ 354.95</b> <del>\$419.80</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
180 capsules	\$ 2.90	<b>\$ 521.95</b> <del>\$629.70</del>	<a href="#">Add to cart</a>

Free AirMail shipping

## Vastarel capsule

### What is this product?

Trimetazidine is indicated in adults as add-on therapy for the symptomatic treatment of patients with stable angina pectoris who are inadequately controlled by or intolerant to first-line antianginal therapies.

### Dosage / Direction for Use

**Dosage:** Oral route.

The dose is one tablet of trimetazidine 35 mg twice daily, i.e. once in the morning and once in the evening, during meals.

The benefit of the treatment should be assessed after three months and trimetazidine should be discontinued if there is no treatment response.

