

Please click price below and you will be redirected to our pharmacy.

Canasa



Generic Canasa is used for treating various forms of mild to moderate inflammation of the colon.

Active ingredient: [Mesalamine](#)

Brand(s): Mesacol Suppository

Manufacturer: Sun Pharmaceutical Industries Ltd.

Disease(s): Ulcerative Colitis

Canasa 500mg

| | Per pill | Per Pack | Order |
|----------------|----------|---|-----------------------------|
| 10 suppository | \$ 3.00 | \$ 29.95 | Add to cart |
| 20 suppository | \$ 3.75 | \$ 74.95 \$74.95 | Add to cart |
| 30 suppository | \$ 3.50 | \$ 104.95 \$104.95 | Add to cart |
| 60 suppository | \$ 2.25 | \$ 134.95 \$179.70 | Add to cart |
| 90 suppository | \$ 2.00 | \$ 179.95 \$269.55 | Add to cart |

Free AirMail shipping

Mesalamine suppository

What is this medicine?

MESALAMINE is used to treat the pain and inflammation caused by ulcerative colitis.

What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions:

- kidney disease
- liver disease
- pyloric stenosis
- an unusual or allergic reaction to mesalamine, salicylates, other medicines, sulfites, foods, dyes, or preservatives
- pregnant or trying to get pregnant
- breast-feeding

How should I use this medicine?

Follow all directions on your prescription label. Do not use this medicine in larger or smaller amounts or for longer than recommended.

Mesalamine rectal comes with patient instructions for using either the rectal suppository or the rectal enema. Follow these directions carefully. Ask your doctor or pharmacist if you have any questions. Do not take mesalamine rectal by mouth. It is for use only in your rectum. Try to empty your bowel and bladder just before using mesalamine rectal. It may be best to use this medicine at bedtime. Wash your hands before and after inserting the rectal suppository. Remove the wrapper before inserting the suppository. Avoid handling the suppository too long or it will melt in your hands. Lie on your back with your knees up toward your chest. Gently insert the suppository into your rectum about 1 inch. For best results, stay lying down for a few minutes. The suppository will melt quickly and you should feel little or no discomfort while holding it in. Avoid using the bathroom for at least 1 hour after using the suppository.

