

## Heximar® Ointment



Heximar® Ointment is a synthetic vitamin D derivative used for the treatment of plaque psoriasis in adults and children above 6 years of age. It lowers the rate with which skin cells grow, thereby controlling the psoriatic condition. It may be used alone or in a combination with other medicines.

Active ingredient: [Calcipotriol](#)

**Brand(s):**Heximar-B

**Manufacturer:** Menarini

**Disease(s):**Plaque Psoriasis

### Heximar® Ointment 15g

	Per pill	Per Pack	Order
1 tube	\$ 44.95	<b>\$ 44.95</b>	<a href="#">Add to cart</a>
2 tube	\$ 39.98	<b>\$ 79.95</b> <del>\$89.90</del>	<a href="#">Add to cart</a>
3 tube	\$ 36.65	<b>\$ 109.95</b> <del>\$134.85</del>	<a href="#">Add to cart</a>
4 tube	\$ 35.99	<b>\$ 143.95</b> <del>\$179.80</del>	<a href="#">Add to cart</a>
5 tube	\$ 34.99	<b>\$ 174.95</b> <del>\$224.75</del>	<a href="#">Add to cart</a>

Free AirMail shipping

## Heximar® Ointment

### What is this product?

Heximar-B® ointment contains 2 complementary active ingredients:

1. calcipotriol – is an analogue of the natural active metabolite of vitamin D3 with a wide range of pharmacological activity in the form of ointment for the topical treatment of psoriasis. It inhibits proliferation of keratinocytes, promotes the differentiation of the epidermis and has a significant effect on cytokines and T-cells in the skin;
  2. betamethasone dipropionate – is a glucocorticosteroid, which has a local anti-inflammatory, antipruritic and immunosuppressive effects.
- ointment affects on all links of pathogenesis of psoriasis and gives a quick effect in the topical treatment of chronic vulgaris plaque psoriasis of any severity;
  - good cosmetic properties (odorless, colorless, does not stain clothes and skin);
  - FDA – approved product.

### Release form:

ointment for external use 15 g

### Composition:

each gram contains:

- calcipotriol anhydrous IP – 50 mcg;
- betamethasone dipropionate equivalent to betamethasone IP – 0.5 mg;
- on the ointment base – q.s.

### Contraindications:

- hypersensitivity to calcipotriol and any auxiliary component of the preparation;
- pregnancy and lactation;
- children up to age 18;
- diseases associated with impaired a calcium metabolism;
- psoriasis of the scalp in the acute stage;
- spotted, erythrodermic, exfoliative and pustular psoriasis;
- renal insufficiency or acute liver failure;
- viral lesions of the skin, fungal or bacterial skin infections, parasitic infections;
- cutaneous manifestations of tuberculosis or syphilis;

