### **Ocuflox**



Generic Ocuflox is used for treating and preventing eye infections associated with conjunctivitis (pink eye) and corneal ulcers caused by certain bacteria in patients 1 year of age and older.

Active ingredient: Ofloxacin

Brand(s):OFLOX Eye Drops / Zenox Eye/Ear Drops Manufacturer: Cipla Limited / Zen Pharma (P) Ltd.

Disease(s):Conjunctivitis

#### Ocuflox 0.3% 5ml

	Per solution	Per Pack	Order
1 solution	\$ 8.00	\$ 8.00	Add to cart
2 solution	\$ 7.00	\$ 14.00 <del>\$16.00</del>	Add to cart
3 solution	\$ 6.50	\$ 19.50 <del>\$24.00</del>	Add to cart
4 solution	\$ 6.00	\$ 24.00 <del>\$32.00</del>	Add to cart
5 solution	\$ 5.50	\$ 27.50 <del>\$40.00</del>	Add to cart
6 solution	\$ 5.00	\$ 30.00 <del>\$48.00</del>	Add to cart

### Ofloxacin 0.3% solution

## What is this medicine?

OFLOXACIN is a quinolone antibiotic. It is used to treat bacterial infections of the eyes.

# What should my health care professional know before I take this medicine?

They need to know if you have any of these conditions:

- an unusual or allergic reaction to ofloxacin, fluoroquinolone antibiotics, other medicines, foods, dyes, or preservatives
- · pregnant or trying to get pregnant
- breast-feeding

# How should I use this medicine?

This medicine is only for use in the eye. Do not take by mouth. Wash hands before and after use. Tilt your head back slightly and pull your lower eyelid down with your index finger to form a pouch. Try not to touch the tip of the dropper to your eye, fingertips, or any other surface. Squeeze the prescribed number of drops into the pouch. Close the eye gently to spread the drops. Your vision may blur for a few minutes. Use your doses at regular intervals. Do not use your medicine more often than directed. Finish the full course prescribed by your doctor or health care professional even if you think your condition is better.

Contact your pediatrician or health care professional regarding the use of this medication in children. Special care may be needed.

Overdosage: If you think you have applied too much of this medicine contact a poison control center or emergency room at once.