

## Mentax



Generic **Mentax** is used for treating certain fungal infections of the skin.

Active ingredient: **Butenafine**

**Brand(s):** Butop

**Manufacturer:** Golden Cross Pharma Pvt. Ltd.

**Disease(s):** Infections

### Mentax 1% 15g

|         | Per tube | Per Pack                              | Order                       |
|---------|----------|---------------------------------------|-----------------------------|
| 1 cream | \$ 14.95 | <b>\$ 14.95</b>                       | <a href="#">Add to cart</a> |
| 2 cream | \$ 12.47 | <b>\$ 24.95</b><br><del>\$29.90</del> | <a href="#">Add to cart</a> |
| 3 cream | \$ 10.98 | <b>\$ 32.95</b><br><del>\$44.85</del> | <a href="#">Add to cart</a> |
| 4 cream | \$ 9.74  | <b>\$ 38.95</b><br><del>\$59.80</del> | <a href="#">Add to cart</a> |
| 5 cream | \$ 8.59  | <b>\$ 42.95</b><br><del>\$74.75</del> | <a href="#">Add to cart</a> |
| 6 cream | \$ 8.33  | <b>\$ 49.95</b><br><del>\$89.70</del> | <a href="#">Add to cart</a> |

### Butenafine 1% cream

#### What is this medicine?

BUTENAFINE is an antifungal medicine. It is used to treat certain kinds of fungal or yeast infections of the skin.

#### What should my health care professional know before I take this medicine?

They need to know if you have any of these conditions:

- an unusual or allergic reaction to butenafine, other medicines, foods, dyes or preservatives
- pregnant or trying to get pregnant
- breast-feeding

#### How should I use this medicine?

This medicine is for external use only. Wash hands before and after use. If treating hand or nail infections, only wash hands before use. Apply a thin layer of the product to cover the affected skin and surrounding area. You can cover the area with a sterile gauze dressing (bandage). Do not use an airtight bandage (such as a plastic-covered bandage). Do not get this medicine in your eyes. If you do, rinse out with plenty of cool tap water. Finish the full course prescribed by your doctor or health care professional even if you think your condition is better. Do not stop using except on your doctor's advice.

Contact your pediatrician or health care professional regarding the use of this medication in children. While this drug may be prescribed for children as young as 12 years of age for selected conditions, precautions do apply.

**Overdosage:** If you think you have applied too much of this medicine contact a poison control center or emergency room at once.

