

Please click price below and you will be redirected to our pharmacy.

## Rulide



Generic **Rulide** is a semi-synthetic macrolide antibiotic. It is used to treat respiratory tract, urinary and soft tissue infections.

Active ingredient: **Roxithromycin**

**Brand(s):**Raxitid

**Manufacturer:** Ranbaxy Laboratories Ltd.

**Disease(s):**Infections

### Rulide 150mg

|            | Per pill | Per Pack                                | Order                       |
|------------|----------|---|-----------------------------|
| 30 tablet  | \$ 1.07  | <b>\$ 31.99</b>                         | <a href="#">Add to cart</a> |
| 60 tablet  | \$ 1.00  | <b>\$ 59.99</b><br><del>\$63.98</del>   | <a href="#">Add to cart</a> |
| 90 tablet  | \$ 0.94  | <b>\$ 84.99</b><br><del>\$95.97</del>   | <a href="#">Add to cart</a> |
| 120 tablet | \$ 0.87  | <b>\$ 104.99</b><br><del>\$127.96</del> | <a href="#">Add to cart</a> |
| 180 tablet | \$ 0.78  | <b>\$ 139.99</b><br><del>\$191.94</del> | <a href="#">Add to cart</a> |
| 360 tablet | \$ 0.69  | <b>\$ 249.99</b><br><del>\$383.88</del> | <a href="#">Add to cart</a> |

Free AirMail shipping

## Roxithromycin tablet

### What is this medicine?

ROXITHROMYCIN is a macrolide antibiotic used for treating various types of infection caused by bacteria, including infections of the top and bottom respiratory tracts, skin and soft tissues, genito-urinary tract (including the infections transferred sexually, except a gonorrhea); spotted fever preventive maintenance at the persons who were in contact to the diseased.

### What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions:

- asthma
- kidney or liver problems
- an unusual or allergic reaction to rulide, other medicines, foods, dyes, or preservatives
- pregnant or trying to get pregnant
- breast-feeding

### How should I use this medicine?

Use Roxithromycin as directed by your doctor. Take the tablet by mouth with food. If you have trouble swallowing the tablet whole, it may be crushed or chewed with a little water.

Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.

Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once.

NOTE: This medicine is only for you. Do not share this medicine with

