

## Adalat CC



Generic Adalat CC Extended-Release is used for treating high blood pressure.

Active ingredient: [Nifedipine](#)

**Brand(s):** Nicardia CD / Nicardia Retard / Nicardia XL

**Manufacturer:** J. B. Chemicals And Pharmaceuticals Ltd

**Disease(s):** Hypertension / Angina

### Adalat CC 30mg

|                       | Per pill | Per Pack                                | Order                       |
|-----------------------|----------|---|-----------------------------|
| 30 tablet             | \$ 3.93  | <b>\$ 118.00</b>                        | <a href="#">Add to cart</a> |
| 60 tablet             | \$ 3.70  | <b>\$ 222.00</b><br><del>\$236.00</del> | <a href="#">Add to cart</a> |
| Free AirMail shipping |          |   |                             |
| 90 tablet             | \$ 3.47  | <b>\$ 312.00</b><br><del>\$354.00</del> | <a href="#">Add to cart</a> |
| Free AirMail shipping |          |   |                             |
| 120 tablet            | \$ 3.35  | <b>\$ 402.00</b><br><del>\$472.00</del> | <a href="#">Add to cart</a> |
| Free AirMail shipping |          |   |                             |

## Nifedipine Extended Release tablet

### What is this medicine?

NIFEDIPINE is a calcium-channel blocker. It affects the amount of calcium found in your heart and muscle cells. This relaxes your blood vessels, which can reduce the amount of work the heart has to do. This medicine is used to treat high blood pressure and chest pain caused by angina.

### What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions:

- heart problems, low blood pressure, slow or irregular heartbeat
- kidney disease
- liver disease
- previous heart attack
- an unusual or allergic reaction to nifedipine, other medicines, foods, dyes, or preservatives
- pregnant or trying to get pregnant
- breast-feeding

### How should I use this medicine?

Take this medicine by mouth with a glass of water. Do not cut, crush or chew. Take your doses at regular intervals. Do not take your medicine more often than directed. Do not suddenly stop taking this medicine. Your doctor will tell you how much medicine to take. If your doctor wants you to stop the medicine, the dose will be slowly lowered over time to avoid any side effects.

Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.

Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once.

