

Please click price below and you will be redirected to our pharmacy.

Ansaid



Ansaid - eye drops, which are used to treat myosis during operations and prevention of the development of reactive postoperative inflammation of the eyes.

Active ingredient: [Flurbiprofen](#)

Brand(s):Ocuflur

Manufacturer: FDC

Disease(s):Myosis / Postoperative Inflammation Of The Eyes.

Ansaid 0.03% 10ml

	Per pill	Per Pack	Order
1 dropper	\$ 24.95	\$ 24.95	Add to cart
2 dropper	\$ 19.98	\$ 39.95 \$49.90	Add to cart
3 dropper	\$ 16.65	\$ 49.95 \$74.85	Add to cart
4 dropper	\$ 14.99	\$ 59.95 \$99.80	Add to cart
5 dropper	\$ 13.99	\$ 69.95 \$124.75	Add to cart

Ansaid 0.03% 5ml

	Per pill	Per Pack	Order
1 dropper	\$ 17.95	\$ 17.95	Add to cart
2 dropper	\$ 12.97	\$ 25.95 \$35.90	Add to cart
3 dropper	\$ 9.98	\$ 29.95 \$53.85	Add to cart
4 dropper	\$ 8.74	\$ 34.95 \$71.80	Add to cart
5 dropper	\$ 7.99	\$ 39.95 \$89.75	Add to cart

Ansaid eye drops

What is Ansaid?

Flurbiprofen a nonsteroidal anti-inflammatory drug (NSAID). This medicine works by reducing hormones that cause inflammation and pain in the body.

Ansaid is used to treat pain or inflammation caused by osteoarthritis or rheumatoid arthritis.

Ansaid may also be used for purposes not listed in this medication guide.

Important Information

Ansaid can increase your risk of fatal heart attack or stroke, especially if you use it long term or take high doses, or if you have heart disease. Do not use Ansaid just before or after heart bypass surgery (coronary artery bypass graft, or CABG).

Ansaid may also cause stomach or intestinal bleeding, which can be fatal. These conditions can occur without warning while you are using this medicine, especially in older adults.

Before taking this medicine

Ansaid can increase your risk of fatal heart attack or stroke, especially if you use it long term or take high doses, or if you have heart disease. Even people without heart disease or risk factors could have a stroke or heart attack while taking this medicine.

Do not use this medicine just before or after heart bypass surgery (coronary artery bypass graft, or CABG).

