

Lariam



Generic Lariam is a medicine used to treat malaria, a disease caused by parasites. This medicine works by interfering with the growth of parasites in the red blood cells of the human body. Parasites that cause malaria typically enter the body through the bite of a mosquito. Malaria is common in areas such as Africa, South America, and Southern Asia. **Generic Lariam** is also used to prevent malaria.

Active ingredient: [Mefloquine](#)

Brand(s):Meflotas

Manufacturer: Intas

Disease(s):Malaria

Lariam 250mg

| | Per pill | Per Pack | Order |
|-----------------------|----------|---|-----------------------------|
| 4 tablet | \$ 9.70 | \$ 38.80 | Add to cart |
| 12 tablet | \$ 9.60 | \$ 115.20 \$116.40 | Add to cart |
| 20 tablet | \$ 9.50 | \$ 190.00 \$194.00 | Add to cart |
| Free AirMail shipping | | | |
| 28 tablet | \$ 9.40 | \$ 263.20 \$271.60 | Add to cart |

Free AirMail shipping

Mefloquine Hydrochloride tablet

What is this medicine?

MEFLOQUINE is used to treat or prevent malaria infections.

What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions:

- depression or history of mental problems including anxiety disorder, schizophrenia, or psychosis
- heart disease
- liver disease
- seizures (epilepsy or convulsions)
- an unusual or allergic reaction to mefloquine, hydroxymefloquine, quinidine, quinine, other medicines, foods, dyes, or preservatives
- pregnant or trying to get pregnant
- breast-feeding

How should I use this medicine?

Take this medicine by mouth with a full glass of water. Take it with food. If you are taking this medicine to prevent malaria, you should start taking it one week before entering the area, and continue for 4 weeks after leaving. Take your doses at regular intervals and on the same day of each week. Do not take your medicine more often than directed. If you are treating an acute malaria infection, you will receive a single dose of the drug. For prolonged travel in an area where malaria is common, consult your healthcare provider for proper dosing schedule.

Talk to your pediatrician regarding the use of this medicine in children. While this drug may be prescribed for children as young as 6 months of age for selected conditions, precautions do apply.

