

## Zanaflex



Generic **Zanaflex** is used for treating muscle spasms. It may also be used to help in the treatment of multiple sclerosis and spinal cord injury.

Active ingredient: [Tizanidine](#)

**Brand(s):** Tizan

**Manufacturer:** Sun Pharmaceutical Industries Ltd.

**Disease(s):** Spasms / Sclerosis

### Zanaflex 2mg

|                       | Per pill | Per Pack                                | Order                       |
|-----------------------|----------|---|-----------------------------|
| 30 tablet             | \$ 1.33  | <b>\$ 39.95</b>                         | <a href="#">Add to cart</a> |
| 60 tablet             | \$ 1.25  | <b>\$ 74.95</b><br><del>\$79.90</del>   | <a href="#">Add to cart</a> |
| 90 tablet             | \$ 1.17  | <b>\$ 104.95</b><br><del>\$119.85</del> | <a href="#">Add to cart</a> |
| 120 tablet            | \$ 1.08  | <b>\$ 129.95</b><br><del>\$159.80</del> | <a href="#">Add to cart</a> |
| 180 tablet            | \$ 1.00  | <b>\$ 179.95</b><br><del>\$239.70</del> | <a href="#">Add to cart</a> |
| Free AirMail shipping |          |   |                             |
| 360 tablet            | \$ 0.97  | <b>\$ 349.95</b><br><del>\$479.40</del> | <a href="#">Add to cart</a> |

Free AirMail shipping

## Tizanidine Hydrochloride tablet

### What is this medicine?

TIZANIDINE helps to relieve muscle spasms. It may be used to help in the treatment of multiple sclerosis and spinal cord injury.

### What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions:

- kidney disease
- liver disease
- low blood pressure
- mental disorder
- an unusual or allergic reaction to tizanidine, other medicines, lactose (tablets only), foods, dyes, or preservatives
- pregnant or trying to get pregnant
- breast-feeding

### How should I use this medicine?

Take this medicine by mouth with a full glass of water. Take this medicine on an empty stomach, at least 30 minutes before or 2 hours after food. Do not take with food unless you talk with your doctor. Take your medicine at regular intervals. Do not take your medicine more often than directed. Do not stop taking except on your doctor's advice. Suddenly stopping the medicine can be very dangerous.

Talk to your pediatrician regarding the use of this medicine in children.

Patients over 65 years old may have a stronger reaction and need a smaller dose.

