Please click price below and you will be redirected to our pharmacy.



Free AirMail shipping

Didanosine Delayed-Release capsule

What is this medicine?

DIDANOSINE, ddl is an antiretroviral drug. It is used with other medicines to treat HIV. This medicine is not a cure for HIV. It will not stop the spread of HIV to others.

What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions:

- gout
- · if you frequently drink alcohol containing drinks
- kidney disease
- liver disease
- pancreatitis
- phenylketonuria
- · tingling or numbness in the hands or feet
- an unusual or allergic reaction to didanosine, other medicines, foods, dyes, or preservatives
- · pregnant or trying to get pregnant
- breast-feeding

How should I use this medicine?

Take this medicine by mouth with a glass of water. Take this medicine on an empty stomach, at least 30 minutes before or 2 hours after food. Do not take with food. Do not chew or crush. Take your medicine at regular intervals. Do not take your medicine more often than directed. For your anti-HIV therapy to work as well as possible, take each dose exactly as prescribed. Do not skip doses or stop your medicine even if you feel better. Skipping doses may make the HIV virus resistant to this medicine and other medicines. Do not stop taking except on your doctor's advice.

Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.

Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once.

NOTE This we distant and for the Denset share this we distant it